

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

described a new genus (*Pseudomachaerium*) of Leguminosae (Dalbergieae) from Paraguay.—P. B. Kennedy (Muhlenbergia 3:8. 1907) has described a new Trifolium from Montana.—T. D. A. Cockerell (*idem* 9) has published a new genus (*Microbahia*) founded on *Syntrichopappus Lemmoni*.—A. A. Heller (*idem* 10–12) has described new western species under Cakile and Ribes.—A. Nelson and P. B. Kennedy (Proc. Biol. Soc. Wash. 19:35–40, 155–158. 1906) have described new species from Nevada under Eriogonum (2), Arabis, Ribes, Gilia, Phlox, Castilleia, Hulsea, Raillardella, Chrysothamnus, Sophia, Sphaerostigma, Godetia, Oreocarya, Cryptanthe (3).—J. M. C.

Ecology and taxonomy of Caulerpa.—SVEDELIUS³⁴ has published the results of his studies of the Ceylonese species of Caulerpa. In a discussion of their mode of life the following subjects are presented: (1) "do all Caulerpas grow under similar external conditions?"; (2) "different ecological types as distinguished by the varying development of their root-system," and (3) "of their assimilation system;" (4) "on the difference between morphological and adaptational characters in Caulerpas." Observations of the different kinds of variations result in the following enumeration: (1) those which depend on locality, and which are to be considered adaptations or "ecologisms;" (2) those which cannot be considered as ecologisms, but are the result of fluctuating variability among the different branchlets; (3) those which can be considered as phylogenetic stages of evolution; (4) bud variations of atavistic origin; (5) dwarf forms; (6) those which do not fall under any of these categories and may be bud variations without atavistic origin (=mutations).

In the taxonomic part of the paper twenty-one species are presented and illustrated, two of them being new.—J. M. C.

Experiments on cell-formation.—By subjecting root tips to various abnormal factors, such as MgSO₄, NaCl, benzine vapor, and extreme temperatures, Němec³ finds that cell walls do not always appear between the daughter nuclei and that the daughter nuclei frequently fuse, the resulting nucleus of course having the double number of chromosomes. In the tips subjected to benzine vapor, the chromosomes are shorter and thicker than the normal ones, although in their formative stages no abnormalities can be observed. He was most successful with root tips of Vicia Faba, Galtonia candicans, Hyacinthus, and various euphorbias. Staminate branches of Larix decidua were placed in a warm chamber and subjected to the action of chloroform vapor. There was little doubt that some pollen mother cells passed directly over into pollen grains. Mixed among these pollen grains true tetrads were occasionally present. Sometimes the pollen mother cells divided once, giving rise to two pollen grains.—W. J. G. Land.

³⁴ SVEDELIUS, NILS, Ecological and systematic studies of the Ceylon species of Caulerpa. Ceylon marine biological reports no. 4. pp. 81–144. *figs.* 51. 1906.

³⁵ N´EMEC, B., Ueber die Bedeutung der Chromosomenzahl. Bull. Internat. Acad. Sci. Bohême 10: pp. 4. 1906.